

PRV

PATENT- OCH REGISTRERINGSVERKET
Patentavdelningen

PCT/ SE 03 / 0 1 0 4 5

Intyg Certificate

Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

REC'D 03 JUL 2003

WIPO

PCT

Ansökan ingavs ursprungligen på engelska.

This is to certify that the annexed is a true copy of the documents as originally filed with the Patent- and Registration Office in connection with the following patent application.

The application was originally filed in English.

(71) Sökande AstraZeneca AB, Södertälje SE
Applicant (s)

(21) Patentansökningsnummer 0201938-8
Patent application number

(86) Ingivningsdatum 2002-06-20
Date of filing

Stockholm, 2003-06-23

För Patent- och registreringsverket
For the Patent- and Registration Office



Sonia André

Avgift
Fee

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)

BEST AVAILABLE COPY

NEW PROCESS

FIELD OF THE INVENTION


The present invention is related to a process for the preparation of 3-cyano-1- naphthoic acid and some analogues thereof, the intermediate 1-bromo-3-cyano naphthalene and some analogues thereof used in this process and a process for the preparation of said intermediate.

BACKGROUND OF THE INVENTION

The compound 3-cyano-1- naphthoic acid and a process for the preparation of the same is previously described in WO 0002859. This process is unattractive for commercial manufacture on account of toxic process effluent arising from use of mercury salt to achieve regioselective decarboxylation, low through process yield and operationally unattractive bromination in concentrated nitric acid.


BRIEF DESCRIPTION OF THE INVENTION

The present invention refers to a process for preparing the compound of formula (1)



wherein X and/or Y₁ and/or Y₂ are independently H, cyano, nitro, trifluoromethoxy, trifluoromethyl, alkoxy, or alkyl and R is H or alkyl either

a) in the case where R=H, by metallo-dehalogenation followed by carboxylation of a compound of formula (12)




wherein X, Y₁ and Y₂ are as defined above, and Hal is Br, I or Cl

or


b) in the case where R=H or alkyl, by palladium mediated carbonylation of a compound of formula (12) with solvolysis.

Furthermore the present invention refers to a compound of formula (12)




wherein X and/or Y₁ and/or Y₂ are independently H, cyano, nitro, trifluoromethoxy, trifluoromethyl, alkoxy, or alkyl and R is H or alkyl and Hal is Br, I or Cl, which is a compound not previously described and which is a key intermediate in the preparation of the compound of formula (1, R=X= Y₁=Y₂ =H) and moreover to a process for preparing a compound of formula (12, Y₁=Y₂=X=H).

Moreover the present invention refers to some other intermediates that may be used in the process for preparing the compound of formula (1, R=X= Y₁=Y₂ =H), namely the compound of formula (20)




the compound of formula (18)




(18)

3-bromo-5-coumalonitrile (27)



(27)


and the compound of formula (63)



(63)

10 DETAILED DESCRIPTION OF THE INVENTION

The process for preparing the compound of formula (1)




(1)

wherein X and/or Y₁ and/or Y₂ are independently H, cyano, nitro, trifluoromethoxy,

15 trifluoromethyl, alkoxy, or alkyl and R is H or alkyl

is carried out either

a) (in the case where R=H) by metallo-dehalogenation followed by carboxylation of a compound of formula (12)




wherein X, Y₁ and Y₂ are as defined above, and Hal is Br, I or Cl

or

- b) (in the case where R=H or alkyl) by palladium mediated carbonylation of a compound
 5 of formula (12) with solvolysis.


Metallo-dehalogenation and carboxylation of compound of formula (12)



10


Metallo-dehalogenation and carboxylation may be carried out by treatment of compound
 (12) with alkyl-lithium reagent, e.g. ⁿBuLi, in THF alone or in admixture with solvents like
 hexane at a temperature below -10 °C, and preferably between -30°C and -75°C, followed
 15 by reaction of the lithiated intermediate with CO₂ and subsequent acidification with e.g.
 HCl.

20

Palladium mediated carbonylation

- 5 The halo-cyano-naphthalene (12) may be reacted with carbon monoxide under elevated pressure, for example between 5 bar and 100 bar, in a solvent such as methanol with an organic base such as triethylamine catalysed by palladium with or without additional phosphine ligand such as triphenyl phosphine or bis-diphenylphosphino propane. The active palladium catalyst can be generated *in situ* from palladium salts such as palladium
- 10 (II) chloride or palladium bis(triphenylphosphine)palladium(II) chloride. The product (1) may be isolated by first of all removing solid residues by filtration and then extracting into aqueous and back into organic with pH control, followed by crystallisation from toluene. The product (6) may be isolated by removing solid residues by filtration followed by crystallisation from solvent.

15

The process for preparing the compound of formula (12, Y₁=Y₂=X=H)(12, Y₁=Y₂=X=H)

wherein Hal is Br, I or Cl

- 20 may be carried out by any of the following routes:

(i) by

(a) treating malic acid (7) with oleum or alternative strongly acid dehydrating media to give coumalic acid (8);

- (b) esterifying coumalic acid (8) to give a pyrone ester (9);
(c) brominating the pyrone ester (9) to give a 3-bromo coumalic ester (10);
(d) reacting the 3-bromo coumalic ester (10) with *in situ* generated benzyne followed by decarboxylation to give a bromonaphthoate (11); and
5 (e) converting/transforming the bromonaphthoate (11) to 1-bromo-3-cyano naphthalene (12, $Y_1=Y_2=X=H$)

or (ii)

by


- 10 (a) treating malic acid (7) with oleum or alternative strongly acid dehydrating media to give coumalic acid (8);

(b) converting coumalic acid (8) into coumalonitrile (25) and subsequently brominating to give 3-bromo-5-coumalonitrile (27); and then
15 (c) converting 3-bromo-5-coumalonitrile (27) into 1-bromo-3-cyano naphthalene (12, $Y_1=Y_2=X=H$)
by cycloaddition of *in situ* generated benzyne, followed by subsequent decarboxylation

or (iii)

20 by

1a) cyanation of 1,2,3,4-tetrahydronaphthalene followed by bromination to give the compound of formula (63)



or

25

1b) bromination of 1,2,3,4-tetrahydronaphthalene followed by cyanodebromination, followed by bromination to give the compound of formula (63); or

1c) bromination of 1,2,3,4-tetrahydronaphthalene followed by metallation and
carboxylation followed by conversion to the 6-cyano-1,2,3,4-tetrahydronaphthalene
followed by bromination to give the compound of formula (63);

5

followed by

2) oxidative aromatization of the compound of formula (63) into 1-bromo-3-cyano
naphthalene (12, $Y_1=Y_2=X=H$);

10 which are illustrated in the reaction schemes below.


15

20

25

30



Route (i)


5

10

Stage (a) - Coumalic acid:


Oleum or alternative strongly acid dehydrating media is added to a suspension of malic acid in a strong acid e.g H_2SO_4 at about $50^\circ C$ to $90^\circ C$, preferably at $75^\circ C$ to $85^\circ C$. Then the mixture is cooled and the product coumalic acid is filtered off.

Coumalic acid (8) is also commercially available.

Stage (b) - Pyrone Ester:

Diisopropylethylamine or other non nucleophilic base (eg DBU) is added to a suspension of coumalic acid in NMP, dimethylsulphate (or else MeBr or MeI) and a non-nucleophilic base, e.g. DBU or iPr_2Net , are added, and the reaction stirred at between $20^\circ C$ and $30^\circ C$.


The reaction mass is diluted, e.g. with toluene, and drowned out into water followed by washing of the organic phase with aqueous bicarbonate and finally water. The solvent is removed by evaporation *in vacuo* and the crude product pyrone ester is purified by filtration isolation from the residual mother liquors.

Stage (c) - 3-Bromo Coumalic Acid:

- 5 Pyrone ester is brominated, e.g. with pyridinium bromide perbromide (pyridinium tribromide) or Br_2 in glacial acetic acid to give 3-bromo coumalic acid .

Stage (d) - Bromonaphthoate:

10




- 15 Isoamyl nitrite and a solution of anthranilic acid in e.g. ethylene glycol dimethyl ether are added to a refluxing solution of a 3-bromo coumalic ester in e.g. ethylene glycol dimethyl ether in the presence of an acid, e.g. catalytic trichloroacetic acid . Benzene-2-diazonium carboxylate is formed by anthranilic acid diazotisation followed by *in situ* decomposition to give benzyne. The reactive benzyne undergoes [4+2] cycloaddition with the 3-bromo coumalic ester to give an intermediate (15) which then extrudes carbon dioxide to give the desired bromonaphthoate. Heating under reflux is continued, the reaction mass is then cooled to about 50°C, a solvent, e.g. toluene is added and the mixture then cooled to ambient. The solution is washed with dilute sodium hydroxide solution , sodium bisulphite solution, water , hydrochloric acid and water again. . The solution is then concentrated *in vacuo* to give the crude bromonaphthoate product.
- 20
- 25

Stage (e) - 1-Bromo-3-Cyano naphthalene :

Three methods for this transformation are possible:

5 Method 1: Conversion to the amide (18) followed by dehydration




10

Bromonaphthoate (11) is heated with ammonia in the presence of a solvent, e.g. toluene, and a catalyst, e.g. KI, at a high temperature to give bromoamide (18)

15 This is followed by dehydration by heating the bromoamide in a large excess of a dehydrating agent, e.g. SOCl₂, to give the compound of formula (12).

Method 2: Conversion to the hydroxamic acid (20)




20

Preparation of hydroxamic acid (20) is achieved by reaction of a bromonaphthoate (11) with hydroxylamine, or a salt thereof, e.g. hydrochloride plus added base. Conversion of the hydroxamic acid (20) to 1-bromo-3-cyano naphthalene (12) is effected by dehydration, e.g. by treatment with PBr₃.


25

Method 3: Direct conversion of Bromonaphthoate (11) to 1-bromo-3-cyano naphthalene (12, X=Y₁=Y₂=H) with Me₂AlNH₂ (21).


- 5 The reagent for this transformation, dimethylaluminium amide, is prepared under strictly anhydrous conditions in an inert atmosphere by condensing anhydrous NH₃ into a solution of AlMe₃ at low temperature.



- 10 A solution of Me₂AlNH₂ solution is added to a solution of a bromonaphthoate in a high-boiling solvent, e.g. m-xylene, and the mixture is heated to reflux. Rapid conversion to the 1-bromo-3-cyano naphthalene (12) occurs and the product is isolated.



15 **Route (ii)**



20

Stage (a) - Coumalic acid: See Route (i) Stage (a) above

Stage (b) - 3-bromo-5-coumalonitrile


Coumalic acid (8) is converted to the corresponding nitrile (25) by conversion to the acid chloride (28) by reaction with a chlorinating agent, e.g. thionyl chloride, followed by reaction with sulfamide ($\text{H}_2\text{NSO}_2\text{NH}_2$).

- 10 Coumalonitrile (25) is brominated using a brominating agent, e.g. pyridinium bromide perbromide (PBPB) in a high-boiling solvent to give bromocoumalonitrile (27). The product is isolated from unreacted starting material by crystallisation.

**Stage (c) - 1-Bromo-3-Cyano naphthalene**


Compound (27) is converted into compound (12) by cycloaddition of *in situ* generated benzyne, followed by subsequent decarboxylation e.g. by heating.

20 The presence of a cyano- rather than an ester group at the 5- position of pyrone ring does not affect the progress of the cycloaddition.



Route (iii)

5




1,2,3,4-tetrahydronaphthalene (also known as Tetralin ®) is cyanated to give cyanotetrahydronaphthalene (70), either directly by reaction with cyanogen bromide with
 10 aluminium chloride as catalyst in carbon disulphide, or via bromotetrahydronaphthalene (68), the resulting cyano tetrahydronaphthalene (70) is brominated to give bromocyanotetrahydronaphthalene (63) which is converted to bromocyanonaphthalene (12) by oxidative aromatisation.

Thus, tetrahydronaphthalene (59) is reacted with bromine, with added iodine as catalyst,
 15 the 6-bromo-1,2,3,4-tetrahydronaphthalene (plus regioisomers) is either a) cyanated by reaction with copper (I) cyanide in NMP at 130 °C for 48h to give 6-cyano-1,2,3,4-tetrahydronaphthalene (70) or b) is lithiated by reaction with n-butyl lithium in THF at -78 °C followed by reaction with carbon dioxide and then dilute hydrochloric acid to furnish 5,6,7,8-tetrahydronaphthalene-2-carboxylic acid (69) along with its regioisomer from
 20 which tetrahydronaphthalene acid (69) is purified by repeated recrystallisation. This acid is converted to cyanonaphthalene (70) by conversion to acid chloride by reaction with thionyl chloride with a small amount of NMP as catalyst, followed by conversion to amide by reaction with ammonia, followed by amide dehydration, for example with PBr_3 .

5,6,7,8-Tetrahydronaphthalene-2-carbonitrile (70) is brominated by reaction with bromine with catalytic ferric bromide in carbon tetrachloride to give bromonitrile (63).

Aromatisation of Substituted Tetralins

5



The aromatisation of the compound of formula (63) into the compound of formula (12) is carried out by heating the compound of formula (63) at a high temperature in the presence of a metal catalyst, e.g. Pd/C. Alternatively, the aromatisation may be carried out for example by stirring with elemental sulphur in a solvent at ambient temperature.

15 PREPARATIONS

Preparation 1

Conversion of malic acid to coumalic acid

Oleum (287g) is added dropwise over 2h to a suspension of malic acid (200g) in concentrated H_2SO_4 (313g) at 75°C and the resulting solution stirred for a further 4h, maintaining temperature at 75°C throughout. The mixture is cooled and then drowned out into ice cold water over 1h. After stirring for 15min and standing overnight, the mixture is cooled to below 10°C and the product is isolated by filtration to give coumalic acid (71 g, 95% purity, 65% yield) after washing and drying.

25 Preparation 2

Conversion of coumalic acid to coumalic acid, methyl ester

Diisopropylethylamine is added to a suspension of coumalic acid (115.5g) in N-methylpyrrolidone (600mL) at 25°C, dimethylsulphate (100.9g) is added over 1h and the reaction stirred at 25°C for 2h. The reaction mass is diluted with toluene, and extracted with water then bicarbonate and finally water. The toluene is removed *in vacuo* and the crude product pyrone ester is purified either by short path distillation or by crystallisation and trituration to give (after removal of residual solvent by evaporation *in vacuo*) the coumalic acid methyl ester (78.8g, 99% purity, 64% yield).

Preparation 3

Conversion of coumalic acid, methyl ester to 3-bromocoumalic acid, methyl ester

A solution of pyrone ester (39g, 95% purity) in acetic acid is added over 3.5hr to a refluxing solution of pyridinium tribromide (105g) in glacial acetic acid (233g). The mixture is held at reflux (85°C -> 107°C) for 3hr then cooled to ambient. Water is added and the crude product is isolated by filtration then washed with water. The crude product is purified by recrystallisation from toluene and *iso*-hexane to give 3-bromocoumalic acid, methyl ester (46g, 82% yield).

Preparation 4

Conversion of 3-bromocoumalic acid, methyl ester to methyl 4-bromo-2-naphthoate

Isoamyl nitrite (24.2g) and a solution of anthranilic acid (28.0g) in ethylene glycol dimethyl ether (90g) are added over 3h to a refluxing solution of 3-bromo coumalic acid, methyl ester (23.3g) in ethylene glycol dimethyl ether (135.8g) in the presence of catalytic trichloroacetic acid (0.165g). The reaction is refluxed for a further 1 hr after the end of addition to ensure complete reaction. The reaction mass is cooled to 50°C, toluene (279g) is added and the mixture then cooled to ambient. The toluene solution is washed with sodium hydroxide solution (75mL, 2M), sodium bisulphite solution (75mL, 5%), water (75mL), hydrochloric acid and water again. The toluene solution is then concentrated *in vacuo* to give methyl 4-bromo-2-naphthoate (30g, 85% purity, 93% yield)..

Preparation 5**Conversion of methyl 4-bromo-2-naphthoate to 4-bromo-2-naphthonitrile**

Dimethylaluminium amide is prepared by the reaction of a solution of trimethylaluminium in toluene (150mL, 2M) with excess anhydrous ammonia (25.5g) at -78°C. Excess ammonia is removed by evaporation at 110 °C and the dimethylaluminium amide solution is then charged to a solution of the bromonaphthoate (39.8g) in m-xylene (321.7g) at 110°C over 1 hour. The reaction is held at 110°C for a further hour and then rapidly cooled to room temperature in ice. The reaction mass is drowned out into aqueous HCl (750 mL, 2M) over 1.5 hours at 5-10°C. The m-xylene solution is concentrated *in vacuo* to give the crude product which is recrystallised from toluene/*iso*-hexane to give 4-bromo-2-naphthonitrile (18.9 g, 54% yield).

Preparation 6**Conversion of methyl 4-bromo-2-naphthoate to 4-bromo-2-naphthonitrile via 4-bromo-2-naphthamide**

To a Carius tube equipped with small magnetic flea and protective outer metal casing is charged methyl 4-bromo-2-naphthoate (1.18 g), aqueous ammonia (9 ml), potassium iodide (0.075 g) and methanol (2 ml). The apparatus is assembled, and lowered into an oil bath at 130 °C. The pressure rises to 4.25 bar. The mixture is heated with stirring under these conditions for 66 h, after which time the assembly is removed from the oil bath and allowed to cool to ambient temperature/pressure. The mixture is cooled to 0 °C to complete crystallisation, and filtered to remove the product. The product is dissolved in EtOAc (50 ml) and washed with 10 % w/v aqueous Na₂CO₃ (2 x 10 ml). The organic layer is separated, dried (MgSO₄) and the solvent removed *in vacuo* to give the product 4-bromo-2-naphthamide as colourless prisms (0.38 g, 94 % str by GC area, 33 % yield,).

To a 10 ml 1-necked round bottomed flask equipped with magnetic stirrer, condenser and inert atmosphere is charged 4-bromo-2-naphthamide (0.093 g) and thionyl chloride (2 ml). The mixture is heated under reflux for 18 h, and the excess thionyl chloride is removed *in vacuo* to afford the crude product 4-bromo-2-naphthonitrile as a yellow solid.

¹H nmr (CDCl₃): 8.15 (s, 1H, ArH), 8.24 (d, 1H, J = 7.4 Hz, ArH), 7.90-7.62 (m, 4H, ArH).

MS: 233 (M⁺), 231 (M⁺), 152, 125, 76.

5

Preparation 7

Conversion of methyl 4-bromo-2-naphthoate to 4-bromo-2-naphthonitrile via 4-bromo-N-hydroxy-2-naphthamide

To a 100 ml 2-necked round bottomed flask equipped with magnetic stirrer, graduated
10 pressure equalised dropping funnel and inert atmosphere is charged bromonaphthoate (2.69 g), hydroxylamine hydrochloride (2.78 g) and methanol (16 ml). 5 M Methanolic KOH (10 ml.) is added dropwise over 40 min to the vigorously stirred suspension at room temperature. An exotherm and an orange colouration is noted on each addition. The reaction mixture (beige suspension) is stirred at room temperature for 17 h after addition of
15 base. The reaction mixture is concentrated to ca. half volume *in vacuo* (water bath < 45 °C) and a 1:1 mixture of water/glacial acetic acid (50 ml) added with vigorous stirring. Stirring is continued for 40 min, and a further portion of 1:1 water/glacial acetic acid (20 ml) added when the suspension becomes too thick to stir. Stirring is continued for 1 h, and the product filtered off under reduced pressure and washed with cold water (3 x 15 ml). The
20 product hydroxamic acid is dried in the vacuum oven at 70 °C to give 4-bromo-N-hydroxy-2-naphthamide as a beige powder (2.2 g, 76 % str. by LC area, 76 % yield,).

To an oven dried 250 ml 2-necked round bottomed flask equipped with magnetic stirrer, condenser, septum and inert atmosphere is charged 4-bromo-N-hydroxy-2-naphthamide (2.0 g) and fluorobenzene (80 ml). Phosphorous tribromide (1.8 ml) is added dropwise
25 over 10 min to the stirred suspension at room temperature and the mixture heated to reflux (85 °C) whereupon a clear orange solution is obtained. Reflux is continued for 18 h, and the solution allowed to cool. The crude reaction mixture is poured into saturated aqueous NaHCO₃ solution (50 ml) and the product extracted with toluene (3 x 50 ml). The combined organic extracts are washed with brine (50 ml) and the solvent removed *in*

vacuo. The residue is crystallised from methanol to give the product 4-bromo-2-naphthonitrile as pale yellow prisms (0.73 g)

The ^1H nmr and mass spectra of the above end product corresponds to those previously
5 obtained.

Preparation 8

Conversion of coumalic acid to 3-bromo-2-oxo-2H-pyran-5-carbonitrile (3-bromocoumalonitrile) via 2-oxo-2H-pyran-5-carbonitrile (coumalonitrile)

10 Coumalic acid (3.91 g) and thionyl chloride (31 ml) are charged to a 100 ml 2-neck round bottomed flask equipped with condenser, magnetic stirrer and inert atmosphere, and the suspension heated to reflux for 1 h. The clear yellow solution is allowed to cool, and the excess thionyl chloride removed *in vacuo*. Sulfamide (3.22 g) is added, and the solid mixture heated to 120°C (bath temp.) for 1 h. The acid chloride melts after a few seconds,
15 and HCl is vigorously evolved. After ca. 15 min, a red foam is obtained, which on further heating collapses to a dark red viscous oil. After 1 h, the reaction mixture has solidified. The reaction mixture is allowed to cool, and transferred to a separating funnel with 10 % w/v aqueous NaHCO_3 solution (150 ml) (heating with the latter being necessary to remove the crude product from the flask). The product is extracted with CH_2Cl_2 (2 x 50 ml) and
20 the combined organic layers washed with sat. NaCl solution (100 ml). The extracts are dried (MgSO_4) and the solvent removed *in vacuo*. The residue is purified by crystallisation from MeOH (2 ml) at 0°C. The product coumalonitrile is obtained as dark orange prisms (1.7 g). Coumalonitrile (2.0g), pyridinium bromide perbromide (5.28g), dimethoxy ethane (13g) and toluene (12,98) are charged to a 100 ml 2-neck round bottomed flask equipped
25 with condenser, magnetic stirrer and inert atmosphere, and heated under reflux for 4 h. The reaction mixture is poured into water (100 ml) and extracted with CH_2Cl_2 (3 x 100 ml). The extracts are dried (MgSO_4) and the solvent removed *in vacuo*. The residue is swirled with ether (20 ml) and the extracts decanted off. The residue is purified by crystallisation from acetone to give the 3-bromo-2-oxo-2H-pyran-5-carbonitrile as an
30 orange powder (1.25 g, 81 % str. by LC area, 31 % yield)

^1H nmr (CDCl_3): 7.74 (d, 1H, J 2.5 Hz, H_a), 8.04 (d, 1H, J = 2.2 Hz, H_b).

MS: 201 (M^+), 199 (M^+), 173, 171, 144, 142, 120, 64, 29.

5 **Preparation 9**

Conversion of 3-bromo-2-oxo-2H-pyran-5-carbonitrile (3-bromocoumalonitrile) to 4-bromo-2-naphthonitrile

Solutions of anthranilic acid in DME (10 ml) and isoamyl nitrite in DME (10 ml, 8.7 g) are added dropwise over 20 min to a stirred solution of 3-bromocoumalonitrile and
10 trichloroacetic acid in DME held at reflux. The mixture is refluxed for a further 10 min, allowed to cool and poured into water (100 ml). The product is extracted with CH_2Cl_2 (2 x 50 ml) and the volatiles removed *in vacuo*. The product crystallises from the residual amyl alcohol at -20°C and the dirty orange solid is collected by filtration *in vacuo*, and dried in the oven at 40°C to give 4-bromo-2-naphthonitrile (0.81 g).

15

Preparation 10

Conversion of 1,2,3,4-Tetrahydronaphthalene to 5,6,7,8-tetrahydronaphthalene-2-carbonitrile

1,2,3,4-Tetrahydronaphthalene (3.3 g), aluminium chloride (6.7 g), cyanogen bromide (5.5
20 g) and carbon disulphide (70 ml) were heated together under reflux for 8 hours however this achieved negligible reaction, the mixture was accordingly concentrated by distilling out solvent at atmospheric pressure until the temperature of the reaction mixture rose to 60°C . Stirring was continued at 60°C for 8 hours, the mixture was cooled, chloroform (100 ml) was added and the resulting mixture then added slowly to a stirred mixture of
25 concentrated hydrochloric acid (3 g) and 50:50 ice water (150 ml) at 0°C . The resulting phases were separated, the aqueous layer was extracted with chloroform (2 x 100 ml), the combined organic phases were washed with saturated aqueous sodium bicarbonate solution (150 ml) and water (2 x 50 ml), they were dried (MgSO_4) and solvent removed by evaporation *in vacuo* to give crude product (3.8 g) comprising a 3:1 mixture of 5,6,7,8-
30 tetrahydronaphthalene-2-carbonitrile and 5,6,7,8-tetrahydronaphthalene-1-carbonitrile.

This was purified by distillation under reduced pressure to give 5,6,7,8-tetrahydronaphthalene-2-carbonitrile in 40% overall yield.

Conversion of 5,6,7,8-tetrahydronaphthalene-2-carbonitrile to 4-bromo-5,6,7,8-tetrahydronaphthalene-2-carbonitrile

5 Bromine (2.5 g, 15.6 mmol) was added cautiously to a stirred mixture of 5,6,7,8-tetrahydronaphthalene-2-carbonitrile (2 g, 12 mmol) and ferric bromide (4.7 g, 15.6 mmol) in carbon tetrachloride (20 ml) at 10 °C. The mixture was stirred at ambient temperature for 8 hours, it was worked up by adding to dilute aqueous hydrochloric acid and extracting
10 with chloroform followed by removal of solvent by evaporation *in vacuo* to give the crude product as a brown oil (5.74 g, 45% purity by gc area, 86% yield). The product was purified by chromatography on silica gel using 1:9 ethyl acetate:hexane eluent to give 4-bromo-5,6,7,8-tetrahydronaphthalene-2-carbonitrile as a mixture of isomers.

15 **Preparation 11**

Conversion of 1,2,3,4-tetrahydronaphthalene to 5-bromo-1,2,3,4-tetrahydronaphthalene and 6-bromo-1,2,3,4-tetrahydronaphthalene

Bromine (66.1 g, 0.41 mol) was added over 3 hours, with stirring at 5 °C to 10 °C, to the 1,2,3,4-tetrahydronaphthalene (50 g, 0.374 mol) along with a small piece of iodine (0.25 g,
20 0.98 mmol). Stirring was continued at ambient temperature for 6 hours and the mixture was then poured slowly into a stirred saturated aqueous solution of sodium sulphite (200 ml) at 10 °C. Stirring was continued for 15 minutes, the resulting mixture was extracted with methylene chloride (3 x 50 ml), the combined organic extracts were washed with water (200 ml), dried (MgSO₄) and solvent removed by evaporation *in vacuo* to give 5-bromo-1,2,3,4-tetrahydronaphthalene along with the 6-bromo-1,2,3,4-tetrahydronaphthalene isomer (86 g, 89.7% purity of combined mono-brominated
25 isomers present in *circa* 3:2 ratio, combined mono-bromo isomer yield 96%).

Conversion of 5-bromo-1,2,3,4-tetrahydronaphthalene and 6-bromo-1,2,3,4-tetrahydronaphthalene to 5,6,7,8-tetrahydronaphthalene-2-carbonitrile along with 5,6,7,8-tetrahydronaphthalene-1-carbonitrile isomer

A mixture of 5-bromo-1,2,3,4-tetrahydronaphthalene and 6-bromo-1,2,3,4-tetrahydronaphthalene (20 g), copper (I) cyanide (8.6 g) and anhydrous N-methylpyrrolidinone (41.3 g) were stirred under dry nitrogen at 130 °C for 40 h. The mixture was cooled to ambient temperature, further N-methylpyrrolidinone (10 g) was added along with saturated aqueous brine (30 ml), the resulting mixture was stirred at ambient for 3 hours and filtered to remove solids. The filtrates were extracted with n-hexane (3 x 50 ml). The combined organic extracts were washed with water (100 ml), dried (MgSO₄) and evaporated *in vacuo* to give crude product (16.2 g). This was purified by distillation to give 5,6,7,8-tetrahydronaphthalene-2-carbonitrile along with regioisomer (13.2 g, 95% purity, 84% yield).

Preparation 12

Conversion of 5-bromo-1,2,3,4-tetrahydronaphthalene and 6-bromo-1,2,3,4-tetrahydronaphthalene to 5,6,7,8-tetrahydronaphthalene-1-carboxylic acid and 5,6,7,8-tetrahydronaphthalene-2-carboxylic acid *n*-Butyl lithium (9.6 ml of 2.5 molar solution in hexane) was added dropwise over 30 minutes to a stirred solution of 5-bromo-1,2,3,4-tetrahydronaphthalene in mixture with its regioisomer 6-bromo-1,2,3,4-tetrahydronaphthalene (5 g) in dry THF (125 ml) and hexane (35 ml) at -70 °C, stirring was continued at -78 °C for 30 minutes, carbon dioxide gas was bubbled through the mixture at -70 °C until no further exotherm was evident, carbon dioxide gas addition was continued for a further 10 minutes as the reaction was allowed to warm to ambient temperature, the mixture was poured into 2 molar aqueous hydrochloric acid (100 ml) and the resulting mixture was extracted with diethyl ether (3 x 50 ml). The combined organic extracts were washed with water (100 ml) and were then extracted with 10% aqueous sodium carbonate solution (3 x 50 ml). The combined aqueous carbonate extracts were acidified carefully by addition of 2 molar hydrochloric acid to adjust the pH to pH 1. The resulting mixture was extracted with diethyl ether (3 x 50 ml), the combined organic

extracts were washed with water (50 ml) and dried (MgSO_4) before solvent was removed by evaporation *in vacuo* to give the crude product in 64% yield comprising a mixture of regioisomers of 5,6,7,8-tetrahydronaphthalene carboxylic acid. This mixture was purified by repeated recrystallisation from ethyl acetate to give 5,6,7,8-tetrahydronaphthalene-2-carboxylic acid as crystallised solid in 93% purity along with 5,6,7,8-tetrahydronaphthalene-1-carboxylic acid as the major component present in the crystallisation mother liquors.

10 **Conversion of 5,6,7,8-tetrahydronaphthalene-2-carboxylic acid to 5,6,7,8-tetrahydronaphthalene-2-carbonitrile**

Acetyl chloride (5 g, 64 mmol) is added dropwise to dry methanol (150 ml) with stirring at ambient temperature under dry nitrogen. Stirring is continued for 15 minutes, 5,6,7,8-Tetrahydronaphthalene-2-carboxylic acid (1 g, 5.7 mmol) is added, the mixture is stirred at ambient temperature for 10 hours and solvent removed by evaporation *in vacuo* to give methyl 5,6,7,8-tetrahydronaphthalene-1-carboxylate. This is then converted to 5,6,7,8-tetrahydronaphthalene-2-carbonitrile using the same procedure described above for conversion of methyl 4-bromo-2-naphthoate to 4-bromo-2-naphthonitrile using dimethylaluminium amide.

20 **Preparation 13**

Conversion of 4-bromo-5,6,7,8-tetrahydronaphthalene-2-carbonitrile to 4-bromo-2-naphthonitrile

4-Bromo-5,6,7,8-tetrahydronaphthalene-2-carbonitrile (0.1 g) was heated with 10% palladium on carbon (1.65 g) under air at 200 °C to 210 °C for 22 hours to give crude 4-bromo-2-naphthonitrile as seen by gc (approximately 75% yield by gc area).

EXAMPLES**Example 1****Conversion of 4-bromo-2-naphthonitrile to 3-cyano-1-naphthoic acid via metallo-dehalogenation and carboxylation.**

- 5 To a 50 ml 4-neck round bottomed flask equipped with a magnetic stirrer, thermometer, septum, CO₂ inlet, N₂ inlet/bubbler and external dry ice/acetone cooling bath is charged (0.35 g, 1.25 mmol), anhydrous hexane (2 ml) and anhydrous THF (8 ml). The suspension is cooled to - 75 °C and BuLi (0.6 ml) added dropwise over 20 min to the vigorously stirred suspension. The bright red solution is stirred for a further 5 min and then carbon
- 10 dioxide bubbled very slowly through the reaction mixture with external cooling. The quenching reaction is very exothermic - maximum temperature reached is - 65 °C. Reaction is judged complete when no further temperature increase is observed upon addition of carbon dioxide. The mixture is stirred at - 65 °C for a further 10 min, and then added cautiously to 2 M HCl. The product is extracted with ethyl acetate (3 x 50 ml), the
- 15 combined extracts dried (MgSO₄) and the solvent removed *in vacuo* to give 3-cyano-1-naphthoic acid.

¹H nmr (D₆DMSO): 7.69-7.87 (m, 2H, 2 x ArH), 8.14 (d, 1H, J = 7.9 Hz, ArH), 8.28 (d, 1H, J = 1.5 Hz, ArH), 8.79 (s, 1H, ArH), 8.85 (d, 1H, J = 8.4 Hz, ArH).

- 20 MS: 197 (M⁺), 180, 152, 125, 29, 18.

Example 2**3-cyano-1-naphthoic acid via carbonylation.**

- Bis(triphenylphosphine)palladium (II) chloride (0.77g) in N-methylpyrrolidinone (170g),
- 25 (10g), triphenyl phosphine (0.57g), and triethylamine (11g,) are mixed in a nitrogen inerted pressure vessel (Parr reactor) at ambient temperature. Water (15.5g) is added and the reactor is repeatedly purged with argon to remove residual air or oxygen. The reactor is vented and then pressurised with carbon monoxide to 7 bar absolute pressure (6 bar gauge pressure) and the mixture stirred at 85 C for 10 hours, maintaining carbon monoxide
- 30 pressure within the reactor at 6 barg. The mixture is cooled to 50 C and vented to

atmospheric pressure, and the reaction mixture then filtered through a bed of celite to remove solids. The filter cake is washed with toluene (160.5g) and then with water (124g). The combined filtrates and washes are allowed to settle and the lower aqueous layer separated. The toluene layer is extracted with water (2 x 124 g). The combined aqueous phase and aqueous extracts were washed with toluene (120g), 2 molar hydrochloric acid (64.5 ml) are added to the aqueous solution over 30 minutes with stirring at 25 to 30 C. The organic layer is separated off and retained, the aqueous layer is extracted with toluene (2 x 120g). The combined organic layer and toluene extracts are mixed with water (62g) and 2 molar sodium hydroxide solution (16.2ml) to extract the product into the aqueous phase. The organic phase is extracted with further water (62g) plus 2 molar sodium hydroxide solution (16.2ml). The combined aqueous extracts are mixed with dichloromethane (350g) and the mixture acidified by addition of 2 molar hydrochloric acid (43ml) over 30 minutes at 25 to 30 C. The lower organic phase is separated and retained; the aqueous phase is extracted with further dichloromethane (100g). The combined dichloromethane solution and extract are washed with 2 molar hydrochloric acid (21.5ml), toluene (120g) is added and dichloromethane is removed by evaporation under reduced pressure to leave a toluene solution of the product. This solution is heated to 60 C, *iso*-hexane (300g) is added over 30 minutes at 60 C, and the mixture cooled over 3 hours to 5 C so as to crystallise the product which is isolated by filtration. The product is washed with pre-cooled *iso*-hexane at 0 C to 5 C and it is then dried overnight in a vacuum oven at 40 C.

The obtained product is confirmed by analysis to be the same as in Example 1.

25 Example 3

Carbonylation of 3-Cyano-1-iodo-2-methoxynaphthalene (12, X=H, Y₁=OMe, Y₂=H, Hal=iodo) to give methyl 3-cyano-2-methoxy-1-naphthoate (1, X=H, Y₁=OMe, Y₂=H)

3-Cyano-1-iodo-2-methoxynaphthalene (10 g, 32.35 mmol), triethylamine (4.91 g, 48.52 mmol), palladium (II) acetate (0.73 g, 3.25 mmol) and methanol (600ml) were charged to a

1000 ml stainless steel Parr Bomb reactor, the vessel was sealed and then repeatedly evacuated and pressurised to 60 psi with carbon monoxide over 4 cycles at ambient temperature so as to completely purge residual air. The reaction mixture was then heated to 75 C with stirring, for 24h, whilst maintaining the carbon monoxide atmosphere at 60 psi pressure. The reactor was then cooled to ambient temperature, the pressure was vented, the reaction mixture was then washed into a receiver vessel using additional methanol and solvent was removed by evaporation *in vacuo*. Solid catalyst residue was removed by dissolving the product into 300ml of methylene chloride and filtering through a short column of silica gel (10 inches depth x 1.5 inches diameter), which was further extracted by additional methylene chloride washes (2 x 500 ml). This crude chromatography step was repeated to further reduce TLC baseline contamination of the product. Solvent was then removed by evaporation *in vacuo* to give the product methyl 3-cyano-2-methoxy-1-naphthoate as a cream coloured solid residue (7.8 g, ca. 100% weight yield). The product identity was confirmed by comparison against reference material by nmr.


The aforementioned procedure was also carried out using 0.2 molar equivalents of palladium (II) acetate catalyst with more reproducible results and also at lower reaction temperature of 60 C over a period of 48 hours, the latter preparation giving the product in 93% yield and 82% purity.

CONCLUSIONS

The new routes described herein offer significantly improved means for large scale manufacture of naphthalene cyanoacid (1) compared with methodology available from the chemical literature. These new routes offer advantage in terms of significantly improved through-route yield (with considerable potential for yet further yield improvement), they avoid the large scale process operability difficulties associated with the previous literature chemistry, they give product of lower cost of manufacture and they avoid the effluent toxicity and reagent toxicity associated with use of stoichiometric mercury salts specified in the previously published chemistry to such products.

CLAIMS


1. A process for preparing the compound of formula (1)



wherein X and/or Y₁ and/or Y₂ are independently H, cyano, nitro, trifluoromethoxy, trifluoromethyl, alkoxy, or alkyl and R is H or alkyl

either

a) in the case where R=H, by metallo-dehalogenation followed by carboxylation of a compound of formula (12)




wherein X, Y₁ and Y₂ are as defined above, and Hal is Br, I or Cl

or

b) (in the case where R=H or alkyl) by palladium mediated carbonylation of a compound of formula (12) with solvolysis.


2. A process for preparing the compound of formula (1) according to claim 1 characterized in that step a) is carried out by treatment of the compound of formula (12) with an alkyl-lithium reagent followed by reaction of the lithiated intermediate with carbon dioxide and then acidification.

3. The compound of formula (12)



wherein X and/or Y₁ and/or Y₂ are independently H, cyano, nitro, trifluoromethoxy, trifluoromethyl, alkoxy, or alkyl and R is H or alkyl and Hal is Br, I or Cl.

- 5 4. A process for preparing a compound of formula (12, Y₁=Y₂=X=H)



(12, Y₁=Y₂=X=H)

wherein Hal is Br, I or Cl

- 10 (i) by

(a) treating malic acid (7) with oleum or alternative strongly acid dehydrating media to give coumalic acid (8);

(b) esterifying coumalic acid (8) to give a pyrone ester (9);

(c) brominating the pyrone ester (9) to give a 3-bromo coumalic ester (10);

- 15 (d) reacting the 3-bromo coumalic ester (10) with *in situ* generated benzyne followed by decarboxylation to give a bromonaphthoate (11); and

(e) converting/transforming bromonaphthoate (11) to 1-bromo-3-cyano naphthalene (12, Y₁=Y₂=X=H)

20

- (ii)

by

(a) treating malic acid (7) with oleum or alternative strongly acid dehydrating media to give coumalic acid (8);

(b) converting coumalic acid (8) into coumalonitrile (25) and subsequently brominating to give 3-bromo-5-coumalonitrile (27); and then

(c) converting 3-bromo-5-coumalonitrile (27) into 1-bromo-3-cyano naphthalene (12,


5 $Y_1=Y_2=X=H$)

by cycloaddition of *in situ* generated benzyne, followed by subsequent decarboxylation

or (iii)

by

10 1a) cyanation of 1,2,3,4-tetrahydronaphthalene followed by bromination to give compound (63)



or

15 1b) bromination of 1,2,3,4-tetrahydronaphthalene followed by cyanodebromination, followed by bromination to give the compound of formula (63); or


1c) bromination of 1,2,3,4-tetrahydronaphthalene followed by carboxylation followed by conversion to the 6-cyano-1,2,3,4-tetrahydronaphthalene followed by bromination to give
20 compound (63); followed by

2) oxidative aromatization of compound (63) into 1-bromo-3-cyano naphthalene (12, $Y_1=Y_2=X=H$).

25 5. A process according to claim 4 characterized in that in process (i) step (d) is carried out by reacting a 3-bromo coumalic ester (10) with *in situ* generated benzyne to give an intermediate (15) followed by decarboxylation to give a bromonaphthoate (11).


6. A process according to claim 4 characterized in that in process (i) step (e) is carried out by either

e1) reaction of compound (11) with ammonia to give compound (18)



followed by dehydration to give compound (12);

e2) reaction of compound (11) with hydroxylamin or a salt thereof to give compound (20);




followed by dehydration to give compound (12);


10 or

e3) direct conversion of compound (11) to compound (12).

7. The compound of formula (20)




8. The compound of formula (18)



20 9. 3-bromo-5-coumalonitrile (27)

10. The compound of formula (63)



100712

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
84

5

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☒ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.